

Name: _____

Date: _____

CHALLENGE!







14.4 Radioactivity

READ 

There are three main types of radiation that involve the decay of the nucleus of an atom:

- **alpha radiation** (α): release of a helium-4 nucleus (two protons and two neutrons). We can represent helium-4 using isotope notation: ${}^4_2\text{He}$. The top number, 4, represents the mass number, and the bottom number represents the atomic number for helium, 2.
- **beta radiation** (β): release of an electron.
- **gamma radiation** (γ): release of an electromagnetic wave.

| | | | |
|---|---|--|---|
|  | Alpha decay | Beta decay | Gamma decay |
| |  |  |  |
| Protons | Decrease by 2 | Increase by 1 | Unchanged |
| Neutrons | Decrease by 2 | Decrease by 1 | Unchanged |

EXAMPLE


Half-life

The time it takes for half of the atoms in a sample to decay is called the half-life. Four kilograms of a certain substance undergo radioactive decay. Let's calculate the amount of substance left over after 1, 2, and 3 half-lives.

- After one half-life, the substance will be reduced by half, to 2 kilograms.
- After two half-lives, the substance will be reduced by another half, to 1 kilogram.
- After three half-lives, the substance will be reduced by another half, to 0.5 kilogram.

So, if we start with a sample of mass m that decays, after a few half-lives, the mass of the sample will be:

| Number of half-lives | Mass left | |
|----------------------|--------------------|-----------------|
| 1 | $\frac{1}{2^1}m =$ | $\frac{1}{2}m$ |
| 2 | $\frac{1}{2^2}m =$ | $\frac{1}{4}m$ |
| 3 | $\frac{1}{2^3}m =$ | $\frac{1}{8}m$ |
| 4 | $\frac{1}{2^4}m =$ | $\frac{1}{16}m$ |



PRACTICE



1. The decay series for uranium-238 and plutonium-240 are listed below. Above each arrow, write "a" for alpha decay or "b" for beta decay to indicate which type of decay took place at each step.
 - a. ${}_{92}^{238}\text{U} \rightarrow {}_{90}^{234}\text{Th} \rightarrow {}_{91}^{234}\text{Pa} \rightarrow {}_{92}^{234}\text{U} \rightarrow {}_{90}^{230}\text{Th} \rightarrow$
 ${}_{88}^{226}\text{Ra} \rightarrow {}_{86}^{222}\text{Rn} \rightarrow {}_{84}^{218}\text{Po} \rightarrow {}_{82}^{214}\text{Pb} \rightarrow {}_{83}^{214}\text{Bi} \rightarrow$
 ${}_{84}^{214}\text{Po} \rightarrow {}_{82}^{210}\text{Pb} \rightarrow {}_{83}^{210}\text{Bi} \rightarrow {}_{84}^{210}\text{Po} \rightarrow {}_{82}^{206}\text{Pb}$
 - b. ${}_{94}^{240}\text{Pu} \rightarrow {}_{95}^{240}\text{Am} \rightarrow {}_{93}^{236}\text{Np} \rightarrow {}_{91}^{232}\text{Pa} \rightarrow {}_{92}^{232}\text{U} \rightarrow$
 ${}_{90}^{228}\text{Bi} \rightarrow {}_{88}^{224}\text{Ra} \rightarrow {}_{89}^{224}\text{Ac} \rightarrow {}_{87}^{220}\text{Fr} \rightarrow {}_{85}^{216}\text{At} \rightarrow$
 ${}_{83}^{212}\text{Bi} \rightarrow {}_{84}^{212}\text{Po} \rightarrow {}_{82}^{208}\text{Pb} \rightarrow {}_{83}^{208}\text{Bi}$
2. Fluorine-18 (${}_{9}^{18}\text{F}$) has a half-life of 110 seconds. This material is used extensively in medicine. The hospital laboratory starts the day at 9 a.m. with 10 grams of ${}_{9}^{18}\text{F}$.
 - a. How many half-lives for fluorine-18 occur in 11 minutes (660 seconds)?
 - b. How much of the 10-gram sample of fluorine-18 would be left after 11 minutes?
3. The isotope ${}_{6}^{14}\text{C}$ has a half-life of 5,730 years. What is the fraction of ${}_{6}^{14}\text{C}$ in a sample with mass, m , after 28,650 years?
4. What is the half-life of this radioactive isotope that decreases to one-fourth its original amount in 18 months?