

Ch. 4 Closure Solutions

CL-4

106 a. $2(y-1)^2 + 8 = 80$

$$\begin{array}{rcl} & -8 & -8 \\ 2(y-1)^2 & = 72 & \\ \text{divide by 2} & & \end{array}$$

$$2(y-1)^2 = 72$$

divide by 2

$$(y-1)^2 = 36$$


✓ ✓

$$y-1 = \pm 6$$

$$\begin{array}{ll} y-1 = 6 & y-1 = -6 \\ \downarrow & \downarrow \\ y = 7 & y = -5 \end{array}$$

c. $y \geq |x+2| - 3$


two variables so solution must be shown on a plane



I'll test $(-2, 0)$

$$0 \geq |-2+2|-3$$

$0 \geq -3$
false



b. $\sqrt{1-2x} = 10$

square both sides

$$1-2x = 100$$

$$-2x = 99$$

$$x = -\frac{99}{2}$$

or -49.5

107

a. $y = \frac{1}{3}x^2 + 1$

$$y = 2x - 2$$

Substitute

$$2x - 2 = \frac{1}{3}x^2 + 1$$

← multiply all terms by 3

$$6x - 6 = x^2 + 3$$

set equal to 0

$$0 = x^2 - 6x + 9$$

$$0 = (x-3)(x-3)$$

$$x-3=0$$

$$x=3$$

$$\begin{array}{l} y = 2(3) - 2 \\ = 4 \end{array}$$

$$(3, 4)$$

A line intersects a parabola at $(3, 4)$ so the line is tangent to the parabola there

107 b

$$y = \sqrt{x-3}$$

$$y = x-5$$

Substitute

$$x-5 = \sqrt{x-3} \quad \leftarrow \text{radical equations can have extraneous solutions so be sure to check answers.}$$

square both sides

↓

$$(x-5)^2 = x-3$$

$$(x-5)(x-5) = x-3$$

$$x^2 - 10x + 25 = x-3$$

↓

$$x^2 - 11x + 28 = 0$$

$$(x-7)(x-4) = 0$$

↓

$$x=7 \quad x=4$$

$$(7)-5 = \sqrt{7-3} \quad 4-5 = \sqrt{4-3}$$

$$2 = \sqrt{4}$$

$$2 = 2$$

✓

$$-1 = \sqrt{1}$$

$$-1 = 1$$

nope

so 4 is extraneous

So, a line intersects a square root function at $(7, 2)$

↑

$$y = 7-5 = 2$$

108

l = cost of lemon pie

B = cost of blueberry pie

two lemon
cost less than 3 less
than 4 blueberry

3 lemon cost 9
more than 3 blueberry

$$2l = 4B - 3$$

$$3l = 3B + 9$$

I'll use substitution!

$$\frac{3l}{3} = \frac{3B+9}{3}$$

$$l = B + 3$$

$$2l = 4B - 3$$

$$2(B+3) = 4B - 3$$

$$\frac{2B+6}{-2B} = \frac{4B-3}{-2B}$$

$$\frac{6}{+3} = \frac{2B-3}{+3}$$

$$2B = 9$$

$$B = \frac{9}{2} = 4.50$$

$$l = 4.50 + 3 \\ = 7.50$$

so Blueberry Pies cost \$4.50 and Lemon cost \$7.50

110a

$$x^2 - 2x - 15 < 0$$

one variable so solution
can be shown on a
number line

Find boundary points

$$x^2 - 2x - 15 = 0$$

$$(x+3)(x-5) = 0$$

$$\downarrow \quad \downarrow$$

$$x = -3 \quad x = 5$$




Test $x=0$

$$0^2 - 2(0) - 15 < 0$$

$$-15 < 0$$

true



(6, 1) (-10, -7)

$$\text{slope} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-7)}{6 - (-10)} = \frac{1 + 7}{6 + 10} = \frac{8}{16} = \frac{1}{2}$$

III
Use (6, 1)

$$y = mx + b$$

$$1 = \frac{1}{2}(6) + b$$

$$\frac{1}{3} = 3 + b$$

$$b = -2$$

110b

$$|3x-2| \geq 10$$

find boundary
points

$$|3x-2| = 10$$

$$3x-2 = 10$$


$$+2 \quad +2$$

$$3x = 12$$

$$3x = -8$$

$$x = 4$$

$$x = -\frac{8}{3}$$




Test $x=0$

$$|3(0)-2| \geq 10$$

$$|-2| \geq 10$$

$2 \geq 10$ false



111a

(6, 1) (-10, -7)

$$\text{slope} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-7)}{6 - (-10)} = \frac{1 + 7}{6 + 10} = \frac{8}{16} = \frac{1}{2}$$

$$y = \frac{1}{2}x - 2$$

112 a

$$2y^2 + 3y = 7$$

$$2y^2 + 3y - 7 = 0$$

use
quad form

$$a=2$$

$$b=3$$

$$c=-7$$

$$x = \frac{-(3) \pm \sqrt{(3)^2 - 4(2)(-7)}}{2(2)}$$

$$x = \frac{-3 \pm \sqrt{65}}{4}$$

112 b

$$3(2x-y) + 12 = 4x - 3$$

$$\begin{array}{r} 6x - 3y + 12 = 4x - 3 \\ -6x \end{array}$$

$$\begin{array}{r} -3y + 12 = -2x - 3 \\ -12 \end{array}$$

$$\begin{array}{r} -3y = -2x - 15 \\ \hline -3 \qquad -3 \qquad -3 \end{array}$$

$$y = \frac{2}{3}x + 5$$

113 b

$$\frac{x^2-9}{x^2+6x+9} \stackrel{\cancel{x^2+9}}{=} \frac{x^2-x-6}{x^2+4} = \frac{(x+3)(x-3)}{(x+3)(x+3)} \stackrel{\cancel{(x+3)}}{=} \frac{(x-3)(x+2)}{x^2+4} = \frac{x-3}{x+3} \cdot \frac{x^2+4}{(x-3)(x+2)}$$

$\cancel{(x+3)}$

$\cancel{(x+3)}$

$\cancel{(x-3)}$

$$= \boxed{\frac{x^2+4}{(x+3)(x+2)}}$$

113 c

$$(6 + \frac{3}{x+1}) \rightarrow \frac{6(x+1)}{(x+1)} + \frac{3}{x+1}$$

common denom
will be $x+1$ condense
to one
fraction

$$\frac{6(x+1) + 3}{x+1} \rightarrow \boxed{\frac{6x+9}{x+1}}$$

113 d

$$\frac{5}{x} - \frac{10}{x^2+2x}$$

factor

$$\rightarrow \frac{5}{x} - \frac{10}{x(x+2)}$$

common denominator
will be $x(x+2)$

$$\frac{5(x+2)}{x(x+2)} - \frac{10}{x(x+2)}$$

condense to a
single fraction

$$\leftarrow \frac{5x+10 - 10}{x(x+2)}$$

$$\frac{5(x+2)}{x(x+2)}$$

$$\boxed{\frac{5}{x+2}}$$

ANSWER

remember
you can't
cancel
anything
out