

1. Define the following terms:

a. Equilibrium

$$\text{net force} = 0$$

b. Force

push or pull, needed to change motion

c. Acceleration

any change in velocity

d. Momentum

mass \times velocity

e. Inertia

tendency of objects to resist changes in motion

f. Friction

force that resists motion

g. Direct relationship between variables

when 1 variable increases, other variable increases

h. Inverse relationship between variables

when 1 variable increases, other decreases

i. Strong relationship between variables

large change in 1 variable results in large change in other

j. Weak relationship between variables

large change in 1 var. results in small change in other

k. Control variable

variable that is constant throughout other

2. What does the 1st Law of Motion state?

objects continue the motion they already have unless an outside acts.

3. What is the mathematical equation related to the 2nd Law of Motion that relates force, mass and acceleration?

$$a = F/m$$

4. What does the 2nd Law of Motion state?

acceleration is directly proportional to force
+ inversely proportional to mass

5. What does the 3rd Law of Motion state?

For EVERY action there is an EQUAL + opposite
REACTION

6. What must be true of the forces acting on objects if their motion is not changing?

net force = 0

7. A train and a car collide. What is true about the forces that each vehicle exerts on the other?

EQUAL strength!

8. In the collision described in the previous question, *why* are the forces as you described?

3rd Law

9. What does the 2nd Law of Motion state about the relationship between force and acceleration?

direct

10. What does the 2nd Law of Motion state about the relationship between mass and acceleration?

inverse

11. In the "1st and 2nd Laws of Motion" lab, what did you data show about the relationship between the force put onto the car and the speed that it attained?

direct

12. In the "1st and 2nd Laws of Motion" lab, what did you data show about the relationship between the mass of the car and the speed that it attained?

inverse

13. In the "1st and 2nd Laws of Motion" lab, why did the car's speed change when the mass increased?

greater inertia, therefore
greater resistance to change in motion

14. What is the SI unit of force?

newton

15. If 3 times the force is applied to the same object, what will be true of its acceleration?

3x greater

16. The action force is "the rifle pushes the bullet forward." What is the reaction force?

bullet pushes rifle back

17. The action force is "the rocket pushes down on the exhaust gases." What is the reaction force?

ex. gases pushes up on rocket

18. An insect and a car windshield collide. If the windshield exerts a 2 N force on the bug, what is the force exerted by the bug on the car windshield?

2N

19. In the "3rd Law of Motion" lab, what was true of the force that moved the cars apart?

EQUAL strength

20. In the "3rd Law of Motion lab, what was true of the speeds of each car when the masses were NOT equal?

speed of larger mass car was lower

21. If a rock has 10 times more mass than an apple, how will the inertia of each object compare?

rock has 10x inertia

22. If a train has more inertia than a bicycle, what will be true about changing the motion of each object?

train has greater resistance to change in motion.

23. What is the mathematical equation that relates momentum, mass and velocity?

$$p = mv$$

24. If a train and a bicycle are moving at the same velocity, but the train has more mass than the bicycle, which has more momentum? Why?

train - more mass

25. If there are 2 identical bicycles, and one has twice the velocity of the other, which one has more momentum? Why?

faster bike - higher veloc.

26. If there were no friction forces at all, and you threw a rock with a 10 N force, how much force would be required to keep it moving at constant velocity?

No force

27. Why is it that we almost never see objects in motion that stay in motion on Earth?

almost always friction!

For each problem below, carry out these steps:

- Write the **formula** that you will use to solve the problem
- Re-write the formula, substituting known values **with units**
- Write the answer using the proper **unit**
- Check your answer for the proper number of **significant figures**
- Check your work for accuracy

1. What is the momentum of a 30.6 kg bicycle moving at 14.2 m/s?

$$p = mv = (30.6 \text{ kg})(14.2 \text{ m/s}) = \underline{435 \text{ kg}\cdot\text{m/s}}$$

2. If a rock has a mass of 18.5 kg and its momentum is 4,200 kg•m/s, what is the velocity of the rock?

$$v = \frac{p}{m} = \frac{4200 \text{ kg}\cdot\text{m/s}}{18.5 \text{ kg}} = \underline{230 \text{ m/s}}$$

3. If a 70. Kg swimmer pushes off the wall with a force of 180 N, what will be the acceleration of the swimmer?

$$a = \frac{F}{m} = \frac{180 \text{ N}}{70. \text{ kg}} = 2.6 \text{ N/kg}$$

4. A construction worker raises a wooden beam with a force of 200. N and accelerates it upward at a rate of 1.3 m/s². What is the mass of the ~~barbell?~~ ^{beam}

$$m = \frac{F}{a} = \frac{200. \text{ N}}{1.3 \text{ m/s}^2} = \underline{150 \text{ kg}}$$

5. How much force is needed to accelerate a 4.0 kg cat at a rate of 5.7 m/s²?

$$F = ma = (4.0 \text{ kg})(5.7 \text{ m/s}^2) = 23 \text{ kg}\cdot\text{m/s}^2 \text{ (N)}$$