1. Define the following terms: a. Energy the ABILITY to to work. b. Work TRANSFER OF ENERGY BY APPLYING FORCE C. Kinetic energy c. Kinetic energy ENERGY OF Moton d. Gravitational potential energy (Stored) EVERGY OF ELEVATED OBJECT e. Elastic potential energy e. Elastic potential energy (STRED) ENERGY OF AN ODJECT HAS BEEN f. Chemical energy f. Chemical energy PE STORED IN MILECHES g. Power RATE OF POING WORK 2. What is the SI unit of work and energy? JOULE (N-m) (kg-m2/2-) 3. What is the original source of practically all of the energy on Earth? .Sunl 4. What does the Law of Conservation of Energy state? ENERGY CAN'T BE CREATED OR DESTROYEL ONLY TRANSFORMED 5. If a force is exerted on an object but it does not move, was work done? Why or why not? NO - No movent = No cork

6. What is the formula for calculating gravitational potential energy? Define each variable. $m = mass(k_3)$ $g = gravity(9.8 n/k_1)$ h=heized VE=mqh 7. If an object has 15 Jof potential energy at a height of 50 cm, how much PE will it have if raised to a height of 100 cm? fim 30 T 8. If the object in the previous question is dropped to the ground, how much kinetic energy will it have at the instant before it hits the ground, assuming air friction is so small that we can ignore it? 9. If one object has 200 J of PE when raised to a certain height, how much PE will a second object have at the same height if the second object has half the mass of the first object? 100] - 1E=mgh 10. An energy car is elevated to the top of a ramp? What is true about the energy of the car? HAS potential ENERGY - GRAVITIONAL 11. The energy car is released from the top of the ramp and rolls down the ramp. What is true about the energy of the car? E IS TRANSFORMING INTO KF 12. The energy car bounces off a rubber band at the bottom of the ramp and rolls back up the ramp? a. How far will the car roll back up the ramp? LOWER HEIGHT THAN RELEASE HEIGHT b. Why will it behave in this way (in terms of energy)? - FRIZITIIN (CHEES + RAMP) CAUSES ENERGY "LOSS" to system TRANSPER OF ENDRU CAR ~ R.B. ~ CAR KE ~ EASTR ~ KE CAUSES "LOSS" to System

13. What is the relationship between the work done to an energy car and the speed that it attains on the track?

KE=/mm



14. A train traveling at 5 m/s has 10,000 J of KE.

a. How much KE will it have if velocity increases to 10 m/s?

10,000 J

b. Does doubling the velocity double the KE? Why or why not?

NO - 4x

For any calculation that you show below, carry out these steps:

- Write the formula that you will use to solve the problem
- Re-write the formula, substituting known values with units
- Write the answer using the proper **unit**
- Check you answer for the proper number of significant figures
- Check you work for accuracy

15. A 25.0 kg bicycle is moving at 7.0 m/s. How much kinetic energy does the bicycle have?

E=1/2 NV ~ =

16. If the bicycle increases its speed to 14. m/s, how much kinetic energy does the bicycle now have?

 $4 \times 610 = 2400$

17. Does doubling the velocity of the bicycle double the kinetic energy of the bicycle? Why or why not? $NO - 4X (v^2)$

18. What happens to the kinetic energy of the bicycle when it is braked to a full stop?

19. If a 12. kg suitcase is lifted to a height of 2.0 m, how much potential energy does the rock now have?

20. If the suitcase is dropped to the ground, how much kinetic energy does the suitcase have in the instant before it hits the floor? (assume air friction is so small that we can ignore it)



21. How much potential energy does the suitcase have when it has fallen halfway to the ground? (again, we can ignore air friction)



22. Fill in the blanks below.

