

1. Define the following terms:

a. Equilibrium

balanced forces, net force = 0

b. Force

pull or push

c. Acceleration

any change in velocity

d. Momentum

strength of motion ($p = mv$)

e. Inertia

property of obj. that resists changes in motion

f. Friction

force that resists motion

g. Direct relationship between variables

increase in 1 var. is related to increase in other

h. Inverse relationship between variables

increase in 1 var. is related to decrease in other var.

i. Strong relationship between variables

Large change in 1 var. is related to large change in other

j. Weak relationship between variables

Large increase in 1 var. is related to small change in other

k. Control variable

• var. kept constant in experiment

2. What does the 1st Law of Motion state?

Objs. tend to keep motion they already have unless a force acts on them

3. What is the mathematical equation related to the 2nd Law of Motion that relates force, mass and acceleration?

$$a = F/m$$

4. What does the 2nd Law of Motion state?acceleration and force are directly proportional
and accel. + mass are inversely proportional

5. What does the 3rd Law of Motion state?

for every action there is an equal + opposite reaction

6. What must be true of the forces acting on objects if their motion is not changing?

balanced, $F_{net} = 0$

7. A train and a car collide. What is true about the forces that each vehicle exerts on the other?

Same!

8. In the collision described in the previous question, why are the forces as you described?

3rd law

9. What does the 2nd Law of Motion state about the relationship between force and acceleration?

directly proportional $a \sim F$

10. What does the 2nd Law of Motion state about the relationship between mass and acceleration?

inversely proportional $a \sim \frac{1}{m}$

11. In the "1st and 2nd Laws of Motion" lab, what did your data show about the relationship between the force put onto the car and the speed that it attained?

more force = more speed

12. In the "1st and 2nd Laws of Motion" lab, what did your data show about the relationship between the mass of the car and the speed that it attained?

more mass = less speed

13. In the "1st and 2nd Laws of Motion" lab, why did the car's speed change when the mass increased?

$$\text{accel} \propto \frac{1}{\text{mass}}$$

14. What is the SI unit of force? newton (N)

15. If 3 times the force is applied to the same object, what will be true of its acceleration?

3x more accel.

16. The action force is "the rifle pushes the bullet forward." What is the reaction force?

the bullet pushes the rifle backward

17. The action force is "the rocket pushes down on the exhaust gases." What is the reaction force?

the exhaust gas pushes up on the rocket

18. An insect and a car windshield collide. If the windshield exerts a 2 N force on the bug, what is the force exerted by the bug on the car windshield?

2 N

19. In the "3rd Law of Motion" lab, what was true of the force that moved the cars apart?

same

20. In the "3rd Law of Motion lab, what was true of the speeds of each car when the masses were NOT equal?

not equal

21. If a rock has 10 times more mass than an apple, how will the inertia of each object compare?

rock has 10x more inertia

22. If a train has more inertia than a bicycle, what will be true about changing the motion of each object?

harder the change motion of train

23. What is the mathematical equation that relates momentum, mass and velocity?

$$p = mv$$

24. If a train and a bicycle are moving at the same velocity, but the train has more mass than the bicycle, which has more momentum? Why?

train, b/c more mass

25. If there are 2 identical bicycles, and one has twice the velocity of the other, which one has more momentum? Why?

bike w/ more velocity

26. If there were no friction forces at all, and you threw a rock with a 10 N force, how much force would be required to keep it moving at constant velocity?

no force

27. Why is it that we almost never see objects in motion that stay in motion on Earth?

friction

28. b/c forces are acting diff. objs.

For each problem below, carry out these steps:

- Write the **formula** that you will use to solve the problem
- Re-write the formula, substituting known values **with units**
- Write the answer using the proper **unit**
- Check you answer for the proper number of **significant figures**
- Check you work for accuracy

1. What is the momentum of a 30.6 kg bicycle moving at 14.2 m/s?

$$p = mv = 30.6 \text{ kg} \cdot 14.2 \text{ m/s} = \frac{434.52}{435} \text{ kg} \cdot \text{m/s}$$

2. If a rock has a mass of 18.5 kg and its momentum is 4,200 kg•m/s, what is the velocity of the rock?

$$v = \frac{p}{m} = \frac{4,200 \text{ kg} \cdot \text{m/s}}{18.5 \text{ kg}} = \frac{227.027\dots}{230} \text{ m/s}$$

3. If a 70. kg swimmer pushes off the wall with a force of 180 N, what will be the acceleration of the swimmer?

$$a = \frac{F}{m} = \frac{180 \text{ N}}{70. \text{ kg}} = \frac{2.5714\dots}{2.6} \text{ N/kg (m/s}^2\text{)}$$

4. A construction worker raises a wooden beam with a force of 200. N and accelerates it upward at a rate of 1.3 m/s². What is the mass of the barbell?

$$m = \frac{F}{a} = \frac{200. \text{ N}}{1.3 \text{ m/s}^2} = \frac{153.84\dots}{150} \text{ kg}$$

5. How much force is needed to accelerate a 4.0 kg cat at a rate of 5.7 m/s²?

$$F = ma = 4.0 \text{ kg} \cdot 5.7 \text{ m/s}^2 = \frac{22.8}{23} \text{ N (kg} \cdot \text{m/s}^2\text{)}$$