


# The Cell Cycle

What controls the life and development of a cell?

**Why?** Most of cell's life is spent in interphase.

An old piece of poetry says "to everything there is a season... a time to be born, a time to die." For cells, the line might say "a time to divide and a time to grow." In multicellular organisms, different types of cells have different roles and need to complete specific tasks. For example, a cell that isn't large enough is not useful for storing nutrients for later, but a cell that is too large will not be useful for transportation through a tiny capillary. In this activity, you will learn about the seasons of a cell's life, and in turn better understand how organisms function.

## Model 1 - The Cell Cycle



1. How many phases are in the cell cycle as shown in the diagram in Model 1?

Four

2. Starting at the starred cell, what is the order of the stages of a cell's life?

G<sub>1</sub>, S, G<sub>2</sub>, M

3. During which phase does the size of the cell increase?

G<sub>1</sub> (Gap 1)

4. During which phase does the number of cells increase?

Mitosis

- 5. Considering your answer to Questions 3 and 4, identify two ways that the growth of an organism can be accomplished through the events of the cell cycle.

Growth can occur during G<sub>1</sub>, when cell size increases or during M<sup>0</sup> (mitosis) when the number of cells increases.



- 6. Cancer, the uncontrolled growth of cells, often results in a tumor, or mass of abnormal cells. Some cancerous tumors consist of many cells that are much smaller than normal. According to Model 1, what part(s) of the cell cycle is (are) most likely being affected?

G<sub>1</sub> - during which cell size increases

- 7. In Model 1, if the length of the arrow represents time, then for those cancerous cells, what happens to the time that is necessary for the cell cycle? What implication might this have for doctors who are treating cancer patients?

If G<sub>1</sub> phase is shortened, then entire cell cycle time is decreased. Cancer cells are produced at a faster rate, so starting treatment ASAP is important.

### Model 2 - Cell Cycle Data

Phase	more Key Process organelles are made	Time Interval (hours)	Sets of DNA present in each cell at end of phase	Number of organelles in each cell at end of phase
Gap <sub>1</sub>	cell growth &	11	1	560
Synthesis	DNA replication	8	2	570
Gap <sub>2</sub>	Protein and organelle synthesis	4	2	600
Mitosis	Cell and nuclear splitting	1	1	300
	Total time:	24		

- 8. Model 2 presents cell cycle data for a typical human cell in culture. Use the phase names in Model 2 to label the G, M, and S phases in Model 1.



- 9. Looking at the third column of Model 2, compare the time spent in mitosis with the time spent in gap<sub>1</sub> in human cells and describe any difference.

More time is spent in Gap 1 than mitosis. (11x as long)

10. Imagine 100 cells were chosen randomly from a tissue sample and examined under a microscope. In which phase of the cell cycle would you expect to find the largest number of cells? Explain.

More cells in gap phase # 1 or synthesis. (part of interphase)



11. Look again at Model 2. Compare the amount of DNA at the beginning and end of synthesis. Why did the amount of DNA change?

Each new cell has to have a full copy of DNA, so DNA is replicated before mitosis.

12. Fill in the "Key Process" column for synthesis phase in Model 2. ✓

13. Cyto = cell, kinesis = cutting. What do you think takes place during cytokinesis?

The cell + cytoplasm are split into two new cells.

14. Other than cytokinesis, what else occurs during the mitosis phase? Hint: Consider the sets of DNA in each cell.

Nuclear division - DNA is evenly divided.


15. Look carefully at information given to you in Model 1 and Model 2. Fill in the key process column in the table for gap<sub>1</sub>. ✓

16. If a culture in the lab starts with one human cell, how many cells will there be after 24 hours?

Two cells.

17. The total time for the phases listed in Model 2 is 24 hours. How many human cells will be in the culture after another 24 hours? Explain.

Four cells



18. Is the original cell "dead" or does it disappear after mitosis? Explain your answer.

No. Original cell is divided into 2 new cells.

19. If a starfish sustains damage to a limb, it often grows a new one. If a human adult sustains damage to his or her spinal cord, mobility is often impaired. If a gecko loses its tail, it may grow a new one. Which type of cell is less likely to go through the cell cycle after being damaged—starfish limb, human spinal cord, or gecko tail? Support your answer.

Human spinal cord cells are less likely so injury remains.

20. Occasionally cells stop dividing and enter another phase, G<sub>0</sub>. If you damage your liver, new liver cells can be produced to replace up to 75% of the liver. However, if you sustain brain damage, your body does not produce new brain cells. Explain this observation using what you have learned about the cell cycle.

Brain cells are in G<sub>0</sub>. If cells are in a G<sub>0</sub> phase then they do not go through cell cycle to divide.